Forest Ecology and Management

Special Issue

Wind and Other Abiotic Risks to Forests

CONTENTS

(Abstracts/contents list published in Biological Abstracts, Biological & Agricultural Index, Current Advances in Ecological Science, Current Awareness in Biological Sciences, Current Contents AB & ES, Ecological Abstracts, Environment Abstracts, Environmental Bibliography, Forestry Abstracts, Geo Abstracts, GEOBASE, Referativnyi Zhurnal)

Introduction 1

Air motion within and above forest vegetation in non-ideal conditions
X. Lee (New Haven, CT, USA) 3

Wind speeds within a single crown of Japanese black pine (Pinus thunbergii Parl.)
J. Zhu, T. Matsuzaki and K. Sakioka (Niigata, Japan) 19

Basic biomechanics of self-supporting plants: wind loads and gravitational loads on a Norway spruce tree
H.-C. Spatz and F. Bruechert (Freiburg, Germany) 33

The mechanics of Norway spruce [Picea abies (L.) Karst]: mechanical properties of standing trees from different thinning regimes
F. Brüchert, G. Becker and T. Speck (Freiburg, Germany) 45

Differences in maximum resistive bending moments of Pinus radiata trees grown on a range of soil types
J.R. Moore (Rotorua, New Zealand) 63

Crown, stem and wood properties of wind-damaged and undamaged Sitka spruce
R.A. Dunham and A.D. Cameron (Aberdeen, UK) 73

An approach for estimating resistance of Japanese cedar to snow accretion damage
A. Kato and H. Nakatani (Tateyama, Japan) 83

Tree breakage from torsional wind loading due to crown asymmetry
S. Skatter and B. Kucera (Aas, Norway) 97

Stem growth responses in Douglas-fir and Sitka spruce following thinning: implications for assessing wind-firmness
S.J. Mitchell (Vancouver, Canada) 105

Irregular heartwood formation in Pinus pinaster Ait. is related to eccentric, radial, stem growth
A. Stokes and S. Berthier (Cestas Gazinet, France) 115

Modelling stem breakage caused by typhoons in plantation Cryptomeria japonica forests
Y. Chiba (Ibaraki, Japan) 123

Wind-induced forces in the near-surface lateral roots of radiata pine
A. Watson (Canterbury, New Zealand) 133

Mechanical stability of Scots pine, Norway spruce and birch: an analysis of tree-pulling experiments in Finland
H. Peltola, S. Kellomäki (Joensuu, Finland), A. Hassinen (Ilomantsi, Finland) and M. Granander (Joensuu, Finland) . . 143

The effects of windthrow on forests at different spatial scales: a review
N.G. Ulanova (Moscow, Russia) 155

Factors influencing windthrow in balsam fir forests: from landscape studies to individual tree studies
J.-C. Ruel (Sainte-Foy, Canada) 169

Frequency and characteristics of severe storms in the Urals and their influence on the development, structure and management of the boreal forests
R. Lässig (Birmensdorf, Switzerland) and S.A. Močalov (Yekaterinburg, Russia) 179

Relationships between climatic parameters and forest vegetation: application to burned area in Alicante (Spain)
F. Belda and J. Meliá (Valencia, Spain) 195

Height growth patterns of Scots pine and Norway spruce in the coastal areas of western Finland
K. Karlsson (Kannus, Finland) 205
Physiological changes in *Pinus sylvestris* needles during early spring under sub-arctic conditions
M.-L. Sutinen (Rovaniemi, Finland), T. Repo (Joensuu, Finland), S. Sutinen (Suonenjoki, Finland), H. Lasarov, L. Alvila and T.T. Pakkanen (Joensuu, Finland) 217
Extent of abiotic damage and its relation to defoliation of conifers in Finland
S. Nevalainen (Joensuu, Finland) and H. Yli-Kojola (Vantaa, Finland) 229
Damage and recovery of tree species after two different tornadoes in the same old growth forest: a comparison of infrequent wind disturbances
C.J. Peterson (Athens, GA, USA) 237
Characterising windthrown gaps from fine spatial resolution remotely sensed data
R.G. Jackson, G.M. Foody (Lewes, UK) and C.P. Quine (Roslin, UK) 253
Management of forests to reduce the risk of abiotic damage — a review with particular reference to the effects of strong winds
B.A. Gardiner and C.P. Quine (Roslin, UK) 261
An investigation of the potential of digital photogrammetry to provide measurements of forest characteristics and abiotic damage
D.R. Miller (Aberdeen, UK), C.P. Quine (Roslin, UK) and W. Hadley (Aberdeen, UK) 279
Models of low temperature and high irradiance and their application to explaining the risk of seedling mortality
K. Blennow (Alnarp, Sweden) and L. Lindkvist (Borås, Sweden) 289
Integration of component models from the tree, stand and regional levels to assess the risk of wind damage at forest margins
A. Talkkari (Hirvas, Finland), H. Peltola, S. Kellomäki and H. Strandman (Joensuu, Finland) 303
Logistic regression models for wind and snow damage in northern Finland based on the National Forest Inventory data
A. Jalkanen (Helsinki, Finland) and U. Mattila (Joensuu, Finland) 315
Using airflow modelling and spatial analysis for defining wind damage risk classification (WINDARC)
V. Lekes and I. Dandul (Olomouc, Czech Republic) 331
A comparison of the relative risk of wind damage to planted forests in Border Forest Park, Great Britain, and the Central North Island, New Zealand
J. Moore (Rotorua, New Zealand) and C.P. Quine (Roslin, UK) 345
A demonstrator of models for assessing wind, snow and fire damage to forests using the WWW
D.R. Miller, R. Dunham, M.L. Broadgate (Aberdeen, UK), R.J. Aspinall (Bozeman, MT, USA) and A.N.R. Law (Aberdeen, UK) 355